

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/339988188>

Using machine learning to predict 1D steady-state temperature profiles from compressible mantle convection simulations

Presentation · November 2019

CITATIONS

0

READS

14

5 authors, including:



Siddhant Agarwal

German Aerospace Center (DLR)

4 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



Nicola Tosi

German Aerospace Center (DLR)

241 PUBLICATIONS 1,120 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



PhD: Unravelling interior evolution of terrestrial planets [View project](#)



Magma ocean solidification and early mantle dynamics [View project](#)

Using machine learning to predict 1-D steady-state temperature profiles from compressible mantle convection simulations

(Towards 1D surrogate modelling)

Siddhant Agarwal, APS DFD 2019

Nicola Tosi

Doris Breuer

Pan Kessel

Grégoire Montavon



Knowledge for Tomorrow



Agenda

- **Introduction** (Mantle Convection and need for Machine Learning)
- **Data and Results I** (using steady-state simulations)
- **Data and Results II** (using evolution simulations)
- **Conclusion**



Introduction




Knowledge for Tomorrow



Introduction


Mantle convection is an important driver of thermal evolution of terrestrial planets



[NASA]



Introduction



- Mantle convection is governed by poorly constrained parameters and initial conditions




Introduction



- Mantle convection is governed by poorly constrained parameters and initial conditions
- In planetary science, the outputs are observable (...sometimes)



Introduction



- Mantle convection is governed by poorly constrained parameters and initial conditions
- In planetary science, the outputs are observable (...sometimes)
- Can we find a mapping using ML to rapidly scan the parameter space?



Data and Results I




Knowledge for Tomorrow




Steady-State Simulations

Traditional 0D surrogate models are limited to simple flows



[Čížková et al. 2017]



[Grott et al. 2011]



Steady-State Simulations

Traditional 0D surrogate models are limited to simple flows

We can leverage ML to find higher-dimensional mappings for more complex flows

Parameters

Ra $\in [1e+4, 1e+9]$ (*vigor of convection*)

RaQ/Ra $\in [0, 10]$ (*vigor of convection from internal heating*)

Di $\in [0, 2]$ (*compressibility*)

η_T $\in [1e+0, 1e+10]$ (*Temperature dependence of viscosity*)

η_V $\in [1e+0, 1e+4]$ (*Pressure dependence of viscosity*)



1-D Temperature profile



Steady-State Simulations

~40000, 2-D, quarter-cylinder, **steady-state** simulations

~9TB of data generated using ~1 million CPU hours

Parameters



$Ra \in [1e+4, 1e+9]$ (*vigor of convection*)

$RaQ/Ra \in [0, 10]$ (*vigor of convection from internal heating*)

$Di \in [0, 2]$ (*compressibility*)

$\eta_T \in [1e+0, 1e+10]$ (*Temperature dependence of viscosity*)

$\eta_v \in [1e+0, 1e+4]$ (*Pressure dependence of viscosity*)




1-D Temperature profile




Steady-State Simulations

~27000 simulations reached statistical steady-state



Steady-State Simulations



Parameters

$Ra \in [1e+4, 1e+9]$ (*vigor of convection*)


$RaQ/Ra \in [0, 10]$ (*vigor of convection from internal heating*)

$Di \in [0, 2]$ (*compressibility*)


$\eta_T \in [1e+0, 1e+10]$ (*Temperature dependence of viscosity*)

$\eta_v \in [1e+0, 1e+4]$ (*Pressure dependence of viscosity*)

Steady-State Simulations




Steady-State Simulations



Steady-State Simulations

Actual —
Predicted - - -



Steady-State Simulations

Average train error: 1.23%


Average test error: 1.25%

Average train error surface heat flux: 6.65%

Average test error surface heat flux: 6.42%


Average train error core heat flux: 22.73%

Average test error core heat flux: 23.54%






Steady-State Simulations

We can now run evolution models




Steady-State Simulations



Steady-State Simulations

Possible sources of error:

- Movement of parameters out of the data manifold during evolution



Steady-State Simulations

Possible sources of error:

- Movement of parameters out of the data manifold during evolution
- Use of the 0D energy equation


$$\rho_m c_p \frac{d}{dt} \int_{R_c}^{R_p} r^2 T(r, t) dr = R_c^2 q_c - R_p^2 q_s + \rho_m \frac{R_p^3 - R_c^3}{3} H_t$$



Steady-State Simulations

Possible sources of error:

- Movement of parameters out of the data manifold during evolution
- Use of the 0D energy equation
- Only 1% of the profiles 'stagnant lid'




Data and Results II



Knowledge for Tomorrow



Evolution Simulations



Parameters

Ra ($\eta_{ref} \in [1e+19, 1e+22 \text{ Pa s}]$)


RaQ ($\Lambda \in [1, 50], \eta_{ref}$)

$T_{initial} \in [1600, 1800 \text{ K}]$

$E \in [1e+5, 5e+5 \text{ J mol}^{-1}]$

$V \in [4e-6, 10e-6 \text{ m}^3 \text{ mol}^{-1}]$


$t=4.5 \text{ Ga}$



Evolution Simulations


Generated ~10000 **evolution** simulations for Mars with:

- Compressible convection (EBA)
- Heat production from core and radiogenic elements
- Temperature and pressure dependent viscosity (Arrhenius)
- Temperature and pressure dependent thermal conductivity and thermal expansion
- Solid phase transitions




Evolution Simulations

Actual —
Predicted - - -




Evolution Simulations



Evolution Simulations

Actual —
Predicted - - -



Conclusion



Knowledge for Tomorrow



Conclusion

- Algorithms like NNs allow for regression in higher-dimensions in space and time
- Data for training on *steady-state* simulations is hard to generate
- Evolution models built from *steady-state* simulations offer reasonable accuracy in certain sub-spaces of the parameter manifold
- Data for training on *evolution* simulations is easier to generate and offers t times more data
- Evolution models learned directly from *evolution* simulations offer better accuracy



Acknowledgements

HPC Resources: North-German Supercomputing Alliance (**HLRN**)

Funding: Helmholtz Einstein International Berlin Research School in Data Science (**HEIBRiDS**)



References

1. Čížková, Hana, et al. "Impact of Compressibility on Heat Transport Characteristics of Large Terrestrial Planets." *Physics of the Earth and Planetary Interiors*, vol. 268, 2017, pp. 65–77., doi:10.1016/j.pepi.2017.04.007
2. Grott, M., et al. "Thermo-Chemical Evolution and Global Contraction of Mercury." *Earth and Planetary Science Letters*, vol. 307, no. 1-2, 2011, pp. 135–146., doi:10.1016/j.epsl.2011.04.040.
3. Hüttig, Christian, and Kai Stemmer. "The Spiral Grid: A New Approach to Discretize the Sphere and Its Application to Mantle Convection." *Geochemistry, Geophysics, Geosystems*, vol. 9, no. 2, 2008, doi:10.1029/2007gc001581.
4. "NASA's InSight Mars Lander." *NASA*, NASA, 26 June 2019, mars.nasa.gov/insight/.
5. "You Can't Do Machine Learning Inside a Database. Can You?" *Datanami*, 28 Mar. 2019, www.datanami.com/2019/04/01/you-cant-do-machine-learning-inside-a-database-can-you/.





Backup Slides





Knowledge for Tomorrow



Steady-State Simulations



Steady-State Simulations



Evolution Simulations

